



In the Claims

Please amend Claims 1, 9, 30, 31 to 33, 46, 54, 66, 74, 86 to 89, 92, 93, and 96, as follows.

1. (Five times amended) A compound of the formula:



wherein,

X is $-O-$, $-S-$, $-NH-$, $[-N(R_2)]$ or $-N-R_2$;

R_2 is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C_3-C_{10}) cycloalkyl, aroyl, (C_2-C_{11}) alkanoyl, and phenylsulfonyl groups;

aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy when p is 2 and X is -O-;

[R₁ is R₂₀, R₂₁ or R₂₂, wherein:

R₂₀ is -(CH₂)_n- where] n is 2, 3, 4, or 5;

[R₂₁ is

-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-,

-CH₂-CH=CH-CH₂-CH₂-,


-CH₂-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-CH₂-, or

-CH₂-CH₂-C≡C-CH₂-,

the -CH=CH- bond being cis or trans;

R₂₂ is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at least one C₁-C₆ linear alkyl group, phenyl group, or




where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂,
-NH₂ or halogen, p is as previously defined;]

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, [monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,]

-C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl, or
 -CH(OR₇)-alkyl[,]; [-C(=W)-alkyl, -C(=W)-aryl, or
 -C(=W)-heteroaryl;]


wherein alkyl is lower alkyl;

aryl is phenyl or



wherein R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy,
 chlorine, fluorine, bromine, iodine, lower
 monoalkylamino, [lower dialkylamino,] nitro, cyano,
 trifluoromethyl, or trifluoromethoxy;

heteroaryl is



Q₃ is -O-, -S-, -NH-, or -CH=N-;

[W is CH₂ or CHR₈ or N-R₉;]

R₇ is hydrogen, lower alkyl, or [alkanoyl] lower alkyl-(C=O)-;

[R₈ is lower alkyl;

R₉ is hydroxy, alkoxy, or -NHR₁₀ ; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above;]

Group Art Unit 1624
Reissue Application No. 09/708,475

November 3, 2004
Attorney Docket No. P25,984-A REI

and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen, [C₁ = 14 C₄] C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁ - C₄ alkoxy, or -COOR₂₃ where R₂₃ is H or C₁ - C₄ alkyl;



with the exclusion of compounds wherein X is -S-, [R₁ is R₂₀,] R is H, and m=1; [all geometric, optical, and stereoisomers thereof,] or a pharmaceutically acceptable acid addition salt thereof.

Group Art Unit 1624
Reissue Application No. 09/708,475

November 3, 2004
Attorney Docket No. P25,984-A REI

9. (Amended Three Times) A compound as claimed in claim 1, wherein X is -O-, -S-, or -NH-; Y is H, Cl, F, or -CF₃; R is selected from the group consisting of hydrogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, -OH, Cl, F, Br, I, [acyl,] C₁-C₃ monoalkylamino, acylamino, [-NO₂-,] -NO₂, -OCF₃, and -CF₃; and n is 2, 3, or 4.

30. (Amended Five Times) A pharmaceutical composition, which comprises a compound of the formula:



wherein X is $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, or $-\text{N}(\text{R}_2)$;

R_2 is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, $(\text{C}_3-\text{C}_{10})$ cycloalkyl, aroyl, $(\text{C}_2-\text{C}_{11})$ alkanoyl, and phenylsulfonyl groups;

wherein aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy when p is 2 and X is $-\text{O}-$;

[R₁ is R₂₀, R₂₁ or R₂₂, wherein:

R₂₀ is $-(CH_2)_n-$ where n is 2, 3, 4, or 5;

[R₂₁ is

$-CH_2-C=CH-CH_2-$,

$-CH_2-C\equiv C-CH_2-$,

$-CH_2-CH=CH-CH_2-CH_2-$,


$-CH_2-CH_2-CH=CH-CH_2-$,

$-CH_2-C\equiv C-CH_2-CH_2-$, or

$-CH_2-CH_2-C\equiv C-CH_2-$,

the $-CH=CH-$ bond being cis or trans;

R₂₂ is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at least one C₁-C₆ linear alkyl group, phenyl group, or




where Z₁ is lower alkyl, $-OH$, lower alkoxy, $-CF_3$, $-NO_2$, $-NH_2$ or halogen, and p as previously defined;]

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, [monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,] $-C(=O)-alkyl$, $-C(=O)-O-alkyl$, $-C(=O)-aryl$, $-C(=O)-heteroaryl$, or $-CH(OR_7)-alkyl$ [,]; $-C(=W)-alkyl$, $-C(=W)-aryl$, or $-C(=W)-heteroaryl$;


alkyl is lower alkyl;

aryl is phenyl or



where R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine, iodine, lower monoalkylamino, [lower dialkylamino,] nitro, cyano, trifluoromethyl, or trifluoromethoxy;

heteroaryl is



Q₃ is -O-, -S-, -NH-, or -CH=N-;

[W is CH₂ or CHR₈ or N-R₉ ;]

R₇ is hydrogen, lower alkyl, or [(C₂-C₁₁) alkanoyl] lower alkyl-(C=O)-;

[R₈ is lower alkyl;

R₉ is hydroxy, alkoxy, or -NHR₁₀ ; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above;] and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is

Group Art Unit 1624

Reissue Application No. 09/708,475

November 3, 2004

Attorney Docket No. P25,984-A REI

hydrogen, C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁ - C₄


alkoxy, or -COOR₂₃ where R₂₃ is H or C₁ - C₄ alkyl;

with the exclusion of compounds wherein X is -S-, [R₁ is R₂₀,] R is H, and m=1;

[all geometric, optical, and stereoisomers thereof,] or a pharmaceutically acceptable

acid addition salt thereof, and a pharmaceutically acceptable carrier therefor.

31. (Amended Five Times) An antipsychotic composition, which comprises a compound of the formula:



wherein

X is $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, or $-\text{N}(\text{R}_2)-$;

R_2 is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, $(\text{C}_3-\text{C}_{10})$ cycloalkyl, aroyl, $(\text{C}_2-\text{C}_{11})$ alkanoyl, and phenylsulfonyl groups; wherein aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy when p is 2 and X is -O-;

[R₁ is R₂₀, R₂₁ or R₂₂, wherein:

R₂₀ is -(CH₂)_n- where] n is 2, 3, 4, or 5;

[R₂₁ is

-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-,

-CH₂-CH=CH-CH₂-CH₂-,


-CH₂-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-CH₂-, or

-CH₂-CH₂-C≡C-CH₂-,

the -CH=CH- bond being cis or trans;

R₂₂ is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at least one C₁-C₆ linear alkyl group, phenyl group, or




where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂,
-NH₂ or halogen, a p is as previously defined;]

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, [monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,] -C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl, or -CH(OR₇)-alkyl[,]; [-C(=W)-alkyl, -C(=W)-aryl, or

-C(=W)-heteroaryl;]


alkyl is lower alkyl;

aryl is phenyl or



where R_5 is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine, iodine, lower monoalkylamino, [lower dialkylamino,] nitro, cyano, trifluoromethyl, or trifluoromethoxy;

heteroaryl is



Q_3 is -O-, -S-, -NH-, or -CH=N-;

[W is CH_2 or CHR_8 or $N-R_9$;]

R_7 is hydrogen, lower alkyl, or [(C_2-C_{11}) alkanoyl] lower alkyl-(C=O)-;

[R_8 is lower alkyl;

R_9 is hydroxy, alkoxy, or -NHR₁₀; and

R_{10} is hydrogen, lower alkyl, C_1-C_3 acyl, aryl,

-C(=O)-aryl or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above;]

and

m is 1, 2, or 3;


with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R

is hydrogen, C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁ - C₄ alkoxy, or -COOR₂₃ where R₂₃ is H or C₁ - C₄ alkyl;

with the exclusion of compounds wherein X is -S-, [R₁ is R₂₀,] R is H, and m=1;

[all geometric, optical, and stereoisomers thereof,] or a pharmaceutically acceptable acid addition salt thereof, in an amount sufficient to produce an antipsychotic effect, and a pharmaceutically acceptable carrier therefor.

32 . (Amended Four Times) A method of treating psychoses, which comprises administering to a mammal a psychoses-treating effective amount of a compound of the formula:



wherein

X is $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, or $-\text{N}(\text{R}_2)-$;

R_2 is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, $(\text{C}_3-\text{C}_{10})$ cycloalkyl, aroyl, $(\text{C}_2-\text{C}_{11})$ alkanoyl, and phenylsulfonyl groups;

where aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower

alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy when p is 2 and X is -O-;

[R₁ is R₂₀, R₂₁ or R₂₂, wherein:

R₂₀ is -(CH₂)_n- where] n is 2, 3, 4, or 5;

[R₂₁ is

-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-,

-CH₂-CH=CH-CH₂-CH₂-,


-CH₂-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-CH₂-, or

-CH₂-CH₂-C≡C-CH₂-,

the -CH=CH- bond being cis or trans;

R₂₂ is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₂ are substituted by at least one C₁-C₆ linear alkyl group, phenyl group, or




where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂,
-NH₂ or halogen, and p is as previously defined;]

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, [monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,] -C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl, or


-CH(OR₇)-alkyl[,]; [-C(=W)-alkyl, -C(=W)-aryl, or
-C(=W)-heteroaryl;]

wherein alkyl is lower alkyl;
aryl is phenyl or



wherein R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy,
chlorine, fluorine, bromine, iodine, lower
monoalkylamino, [lower dialkylamino,] nitro, cyano,
trifluoromethyl, or trifluoromethoxy;

heteroaryl is



Q₃ is -O-, -S-, -NH-, or -CH=N-;

[W is CH₂ or CHR₈ or N-R₉;]

R₇ is hydrogen, lower alkyl, or [(C₂-C₁₁) alkanoyl] lower alkyl-(C=O)-;

[R₈ is lower alkyl;

R₉ is hydroxy, alkoxy, or -NHR₁₀; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above;]

and


m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen, C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃

wherein R₂₃ is H or C₁-C₄ alkyl;

with the exclusion of compounds wherein X is -S-, [R₁ is R₂₀,] R is H, and m=1; [all geometric, optical, and stereoisomers thereof,] or a pharmaceutically acceptable acid addition salt thereof.

33. (Amended Four Times) An analgesic composition, which comprises a compound of the formula:



wherein,

X is $-O-$, $-S-$, $-NH-$, or $[-N(R_2)] \overset{|}{-N(R_2)}$;

R_2 is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C_3-C_{10}) cycloalkyl, aroyl, (C_2-C_{11}) alkanoyl, and phenylsulfonyl groups;

wherein aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy when p is 2 and X is -O-;

[R₁ is R₂₀, R₂₁ or R₂₂, wherein:

R₂₀ is -(CH₂)_n- where] n is 2, 3, 4, or 5;

[R₂₁ is

-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-,

-CH₂-CH=CH-CH₂-CH₂-,


-CH₂-CH₂-CH=CH-CH₂-,

-CH₂-C≡C-CH₂-CH₂-, or

-CH₂-CH₂-C≡C-CH₂-,

the -CH=CH- bond being cis or trans;

R₂₂ is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at least one C₁-C₆ linear alkyl group, phenyl group, or




where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂, -NH₂ or halogen, and p is as previously defined;]

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, [monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,] -C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl, or

$-\text{CH}(\text{OR}_7)-\text{alkyl}[,]; [-\text{C}(=\text{W})-\text{alkyl}, -\text{C}(=\text{W})-\text{aryl}, \text{ or } -\text{C}(=\text{W})-\text{heteroaryl};]$


wherein alkyl is lower alkyl;

aryl is phenyl or



wherein R_5 is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine, iodine, lower monoalkylamino, [lower dialkylamino,] nitro, cyano, trifluoromethyl, or trifluoromethoxy;

heteroaryl is



wherein Q_3 is $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, or $-\text{CH}=\text{N}-$;

[W is CH_2 or CHR_8 or $\text{N}-\text{R}_9$;]

R_7 is hydrogen, lower alkyl, or [(C_2-C_{11}) alkanoyl] lower alkyl-($\text{C}=\text{O}$)-;

[R_8 is lower alkyl;

R_9 is hydroxy, alkoxy, or $-\text{NHR}_{10}$; and

R_{10} is hydrogen, lower alkyl, C_1-C_3 acyl, aryl,

$-\text{C}(=\text{O})-\text{aryl}$ or $-\text{C}(=\text{O})-\text{heteroaryl}$,

where aryl and heteroaryl are as defined above;] and

m is 1, 2, or 3;

Group Art Unit 1624
Reissue Application No. 09/708,475


November 3, 2004
Attorney Docket No. P25,984-A REI

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen, C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃

wherein R₂₃ is H or C₁-C₄ alkyl;

with the exclusion of compounds wherein X is -S-, [R₁ is R₂₀,] R is H, and m=1; [all geometric, optical, and stereoisomers thereof,] or a pharmaceutically acceptable acid addition salt thereof, in an amount sufficient to produce a pain-relieving effect, and a pharmaceutically acceptable carrier therefor.

46. (Amended Twice) A compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-C₁₀) cycloalkyl, aroyl, (C₂-C₁₁) alkanoyl, and phenylsulfonyl groups;

aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is


-CH₂-CH=CH-CH₂-,
-CH₂-C≡C-CH₂-,
-CH₂-CH=CH-CH₂-CH₂-,
-CH₂-CH₂-CH=CH-CH₂-,
-CH₂-C≡C-CH₂-CH₂-, or
-CH₂-CH₂-C≡C-CH₂-,

the -CH=CH- bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, dialkylaminocarbonyl, formyl, -C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl, -CH(OR₇)-alkyl, -C(=W)-alkyl, -C(=W)-aryl, or -C(=W)-heteroaryl;


wherein alkyl is lower alkyl;

aryl is phenyl or



wherein R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine, iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl, or trifluoromethoxy;

heteroaryl is



wherein Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂ or CHR₈ or N-R₉;

R₇ is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl, -C(=O)-aryl, or
-C(=O)-heteroaryl,

wherein aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;


with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is
hydrogen, C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄
alkoxy, or -COOR₂₃.

wherein R₂₃ is H or C₁-C₄ alkyl;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable
acid addition salt thereof.

54. (Amended Twice) A compound as claimed in claim 46, wherein X is -O-, -S-, or -NH-; Y is H, Cl, F, or -CF₃; R is selected from the group consisting of hydrogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, -OH, Cl, F, Br, I, C₁-C₃ monoalkylamino, acylamino, -NO₂, -OCF₃, and -CF₃.

66. (Amended Twice) A compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-C₁₀) cycloalkyl, aroyl, (C₂-C₁₁) alkanoyl, and phenylsulfonyl groups;


aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at least one C₁-C₆ linear alkyl group, phenyl group or



wherein Z_1 is lower alkyl, $-\text{OH}$, lower alkoxy, $-\text{CF}_3$, $-\text{NO}_2$, $-\text{NH}_2$ or halogen;

R_{20} is $-(\text{CH}_2)_n-$, where n is 2, 3, 4 or 5;

R_{21} is

$-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-$,

$-\text{CH}_2-\text{C}\equiv\text{C}-\text{CH}_2-$,

$-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}_2-$,

$-\text{CH}_2-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-$,

$-\text{CH}_2-\text{C}\equiv\text{C}-\text{CH}_2-\text{CH}_2-$, or

$-\text{CH}_2-\text{CH}_2-\text{C}\equiv\text{C}-\text{CH}_2-$,

the $-\text{CH}=\text{CH}-$ bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine,

fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro,

lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl,

trifluoroacetyl, aminocarbonyl, monoalkylaminocarbonyl,


dialkylaminocarbonyl, formyl, $-\text{C}(=\text{O})$ -alkyl, $-\text{C}(=\text{O})$ -O-alkyl,

$-\text{C}(=\text{O})$ -aryl, $-\text{C}(=\text{O})$ -heteroaryl, $-\text{CH}(\text{OR}_7)$ -alkyl, $-\text{C}(=\text{W})$ -alkyl,

-C(=W)-aryl, or -C(=W)-heteroaryl;


wherein alkyl is lower alkyl;

aryl is phenyl or



wherein R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy,
chlorine, fluorine, bromine, iodine, lower
monoalkylamino, lower dialkylamino, nitro, cyano,
trifluoromethyl, or trifluoromethoxy;

heteroaryl is



wherein Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂ or CHR₈ or N-R₉;

R₇ is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and


R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

wherein aryl and heteroaryl are as defined above; and
m is 1, 2, or 3;
with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is
hydrogen, C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄
alkoxy, or -COOR₂₃
wherein R₂₃ is H or C₁-C₄ alkyl;
with the exclusion of compounds wherein X is -S-, R₁ is R₂₀, R is H, and m=1;
all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable
acid addition salt thereof.

74. (Amended Twice) A compound as claimed in claim 66, wherein X is -O-, -S-, or -NH-; Y is H, Cl, F, or -CF₃; R is selected from the group consisting of hydrogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, -OH, Cl, F, Br, I, C₁-C₃ monoalkylamino, acylamino, -NO₂, -OCF₃, and -CF₃; and n is 2, 3, or 4.

86. (Amended) A pharmaceutical composition, which compromises a compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-C₁₀)cycloalkyl, aroyl, (C₂-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;


Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is

-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-;

-CH₂-CH=CH-CH₂-CH₂-;



the -CH=CH- bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine,

bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl

thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl,


aminocarbonyl, dialkylaminocarbonyl, formyl,

-C(=O)-alkyl , -C(=O)-O-alkyl , -C(=O)-aryl , -C(=O)-heteroaryl ,

$\text{-CH(OR}_7\text{)-alkyl}$, -C(=W)-alkyl , -C(=W)-aryl , or -C(=W)-heteroaryl ;

where alkyl is lower alkyl;

aryl is phenyl or




where R_5 is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine,

iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl,

trifluoromethoxy;

heteroaryl is



where Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂ or CHR₈ or N-R₉;

R₇ is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,


C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃,

where R₂₃ is C₁-C₄ alkyl;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid

addition salt thereof, and a pharmaceutically acceptable carrier therefor.

87. (Amended) A pharmaceutical composition, which comprises a compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-

C₁₀)cycloalkyl, aroyl, (C₂-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is defined hereinafter;

p is 1 or 2;


Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy,

trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at

least one C₁-C₆ linear alkyl group, phenyl group or



where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂, -NH₂ or halogen;

R₂₀ is -(CH₂)_n-, where n is 2, 3, 4 or 5;

R₂₁ is

-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-;

-CH₂-CH=CH-CH₂-CH₂-;

-CH₂-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-CH₂-; or

-CH₂-CH₂-C≡C-CH₂-;

the -CH=CH- bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine,

iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio,

trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl,


aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,

-C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl,

-CH(OR₇)-alkyl, -C(=W)-alkyl, -C(=W)-aryl, or -C(=W)-heteroaryl;


where alkyl is lower alkyl;

aryl is phenyl or



where R_5 is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine,
iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl,
trifluoromethoxy;

heteroaryl is



where Q_3 is -O-, -S-, -NH-, or -CH=N-;

W is CH_2 , or CHR_8 or $N-R_9$;

R_7 is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R_8 is lower alkyl;

R_9 is hydroxy, lower alkoxy, or -NHR₁₀; and

R_{10} is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,

C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂,


where R₂ is C₁-C₄ alkyl;

with the exclusion of compounds wherein X is -S-, R₁ is R₂₀, R is H, and m=1;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid

addition salt thereof, and a pharmaceutically acceptable carrier therefor.

88. (Amended) An antipsychotic composition, which comprises a compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-

C₁₀)cycloalkyl, aroyl, (C₂-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is

-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-;

-CH₂-CH=CH-CH₂-CH₂-;

-CH₂-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-CH₂-; or


-CH₂-CH₂-C≡C-CH₂-;

the -CH=CH- bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, dialkylaminocarbonyl, formyl,
-C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl,
-CH(OR₇)-alkyl, -C(=W)-alkyl, -C(=W)-aryl, or -C(=W)-heteroaryl;


where alkyl is lower alkyl;

aryl is phenyl or



where R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine, iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl, trifluoromethoxy;

heteroaryl is



where Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂, or CHR₈, or N-R₉;

R₇ is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;


with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,

C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃

where R₂₃ is C₁-C₄ alkyl;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid addition salt thereof, in an amount sufficient to produce an antipsychotic effect, and a pharmaceutically acceptable carrier therefor.

89. (Amended) An antipsychotic composition, which comprises a compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-C₁₀)cycloalkyl, aroyl, (C₂-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is defined hereinafter;


p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is R₂₀ or R₂₁ in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at

least one C₁-C₆ linear alkyl group, phenyl group or



where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂, -NH₂ or halogen;

R₂₀ is -(CH₂)_n-, where n is 2, 3, 4 or 5;

R₂₁ is

-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-;

-CH₂-CH=CH-CH₂-CH₂-;

-CH₂-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-CH₂-; or

-CH₂-CH₂-C≡C-CH₂-;

the -CH=CH- bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine,

iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio,

trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl,


aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,

-C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl,

-CH(OR₇)-alkyl, -C(=W)-alkyl, -C(=W)-aryl, or -C(=W)-heteroaryl;


where alkyl is lower alkyl;

aryl is phenyl or



where R_5 is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine, iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl, trifluoromethoxy;

heteroaryl is



where Q_3 is -O-, -S-, -NH-, or -CH=N-;

W is CH_2 or CHR_8 or $N-R_9$;

R_7 is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R_8 is lower alkyl;

R_9 is hydroxy, lower alkoxy, or -NHR_{10i}; and

R_{10} is hydrogen, lower alkyl, C_1-C_3 acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,

C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂,

where R₂ is C₁-C₄ alkyl;


with the exclusion of compounds wherein X is -S-, R₁ is R₂₀, R is H, and m=1;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid

addition salt thereof, in an amount sufficient to produce an antipsychotic effect,

and a pharmaceutically acceptable carrier therefor.

92. (Amended) An analgesic composition, which comprises a compound of the formula



wherein

$$\begin{array}{c} | \\ \text{X is -O-, -S-, -NH-, or -N(R}_1\text{);} \end{array}$$

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-C₁₀)cycloalkyl, aroyl, (C₇-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy, trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is

-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-;

-CH₂-CH=CH-CH₂-CH₂-;

-CH₂-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-CH₂-; or

-CH₂-CH₂-C≡C-CH₂-;

the -CH=CH- bond being cis or trans;


R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio, trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl, aminocarbonyl, dialkylaminocarbonyl, formyl,

-C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl,

-CH(OR₇)-alkyl, -C(=W)-alkyl, -C(=W)-aryl, or -C(=W)-heteroaryl;

where alkyl is lower alkyl;

aryl is phenyl or




where R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine,

iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl,

trifluoromethoxy;

heteroaryl is



where Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂ or CHR₈ or N-R₉;

R₇ is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,

C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃


where R₂₃ is C₁-C₄ alkyl;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid

addition salt thereof, in an amount sufficient to produce a pain-relieving effect,

and a pharmaceutically acceptable carrier therefor.

93. (Amended) An analgesic composition, which comprises a compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-

C₁₀)cycloalkyl, aroyl, (C₇-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is defined hereinafter;

p is 1 or 2;


Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy,

trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

(R₁) is R₂₀ or R₂₁, in which one or more carbon atoms of R₂₀ or R₂₁ are substituted by at

least one C₁-C₆ linear alkyl group, phenyl group or



where Z₁ is lower alkyl, -OH, lower alkoxy, -CF₃, -NO₂, -NH₂ or halogen;

R₂₀ is -(CH₂)_n-, where n is 2, 3, 4 or 5;

R₂₁ is

-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-;

-CH₂-CH=CH-CH₂-CH₂-;

-CH₂-CH₂-CH=CH-CH₂-;

-CH₂-C≡C-CH₂-CH₂-, or

-CH₂-CH₂-C≡C-CH₂-;

the -CH=CH- bond being cis or trans;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine, bromine,

iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio,

trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl,


aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, formyl,

-C(=O)-alkyl, -C(=O)-O-alkyl, -C(=O)-aryl, -C(=O)-heteroaryl,

-CH(OR₇)-alkyl, -C(=W)-alkyl, -C(=W)-aryl, or -C(=W)-heteroaryl;

where alkyl is lower alkyl;

aryl is phenyl or




where R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine,

iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano, trifluoromethyl,

trifluoromethoxy;

heteroaryl is



where Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂ or CHR₈ or N-R₉;

Group Art Unit 1624
Reissue Application No. 09/708,475

November 3, 2004
Attorney Docket No. P25,984-A REI

R₇ is hydrogen, lower alkyl, or acyl;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl, or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,

C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃,

where R₂₃ is C₁-C₄ alkyl;


with the exclusion of compounds wherein X is -S-, R₁ is R₂₀, R is H, and m=1;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid

addition salt thereof, in an amount sufficient to produce a pain-relieving effect,

and a pharmaceutically acceptable carrier therefor.

96. (Amended) A compound of the formula



wherein

X is -O-, -S-, -NH-, or -N(R₂);

R₂ is selected from the group consisting of lower alkyl, aryl lower alkyl, aryl, (C₃-

C₁₀)cycloalkyl, aroyl, (C₂-C₁₁)alkanoyl, and phenylsulfonyl groups;

aryl is as defined hereinafter;

p is 1 or 2;

Y is hydrogen, lower alkyl, hydroxy, chlorine, fluorine, bromine, iodine, lower alkoxy,

trifluoromethyl, nitro, or amino, when p is 1;

Y is lower alkoxy, hydroxy, or halogen when p is 2 and X is -O-;

n is 2, 3, 4 or 5;

R is hydrogen, lower alkyl, lower alkoxy, hydroxyl, carboxyl, chlorine, fluorine,

bromine, iodine, amino, lower mono or dialkylamino, nitro, lower alkyl thio,

trifluoromethoxy, cyano, acylamino, trifluoromethyl, trifluoroacetyl,


aminocarbonyl, dialkylaminocarbonyl, formyl, -C(=O)-alkyl, -C(=O)-O-alkyl,

-C(=O)-aryl, -C(=O)-heteroaryl, or -CH(OR₇)-alkyl; -C(=W)-alkyl,

-C(=W)-aryl, or -C(=W)-heteroaryl;

alkyl is lower alkyl;

aryl is phenyl or




where R₅ is hydrogen, lower alkyl, lower alkoxy, hydroxy, chlorine, fluorine, bromine,

iodine, lower monoalkylamino, lower dialkylamino, nitro, cyano,

trifluoromethyl, or trifluoromethoxy;

heteroaryl is



Q₃ is -O-, -S-, -NH-, or -CH=N-;

W is CH₂ or CHR₈ or N-R₉;

R₇ is hydrogen, lower alkyl, or lower alkyl-(C=O)-;

R₈ is lower alkyl;

R₉ is hydroxy, lower alkoxy, or -NHR₁₀; and

Group Art Unit 1624
Reissue Application No. 09/708,475

November 3, 2004
Attorney Docket No. P25,984-A REI

R₁₀ is hydrogen, lower alkyl, C₁-C₃ acyl, aryl,

-C(=O)-aryl or -C(=O)-heteroaryl,

where aryl and heteroaryl are as defined above; and

m is 1, 2, or 3;

with the proviso that at least one R is selected from the group consisting of

dialkylaminocarbonyl, formyl, -C(=W)-alkyl, -C(=W)-aryl, and

-C(=W)-heteroaryl;

with the exclusion of compounds wherein X is O or S, Y is hydrogen, and R is hydrogen,

C₁-C₄ alkyl, chlorine, fluorine, bromine, iodine, cyano, C₁-C₄ alkoxy, or -COOR₂₃

where R₂₃ is C₁-C₄ alkyl;

with the exclusion of compounds wherein X is -S-, R is H, and m=1;

all geometric, optical and stereoisomers thereof, or a pharmaceutically acceptable acid

addition salt thereof.